

Equally Well

AOTEAROA, NEW ZEALAND - SOME KEY FACTS




Life expectancy reduced by up to 25 years

Two-thirds of premature mortality for people experiencing mental health and addiction issues is due to **preventable and treatable** physical health conditions - particularly cancers and cardiovascular disease.

Co-existing issues are the norm

Around **3 in 5 adults** experiencing mental health issues report having one or more long-term physical health issues.



Cancer

The likelihood of cancer survival for people who experience mental health and addiction issues is reduced by **3 times** for colorectal cancer and **2.5 times** for breast cancer.

Medication

Some psychiatric and addiction treatment medications contribute to poorer health outcomes, for example cardiometabolic issues, particularly weight gain, and poor oral health.



Substance use

People with problematic alcohol use have almost **double the risk** of developing a wide range of cancers.

People with methamphetamine use disorders are more likely to experience heart disease, cerebrovascular complications, oral health diseases and increased transmission of blood borne viruses.

75% of people who receive opioid substitution treatment are likely to have Hepatitis C.



1. Cunningham, R., Sarfati, D., Peterson, D., Stanley, J. & Collings, S. (2014). *Premature mortality in adults using New Zealand psychiatric services*. New Zealand medical journal, 127(1394), 31-41.
2. Scott, K. M., Bruffaerts, R., Simon, G. E., Alonso, J., Angermeyer, M., de Girolamo, G., Von Korff, M. (2007). *Obesity and mental disorders in the general population: Results from the world mental health surveys*. International Journal of Obesity, 32(1), 192-200. doi:10.1038/sj.ijo.0803701.
3. Cunningham, R., Sarfati, D., Stanley, J., Peterson, D. & Collings, S. (2015). *Cancer survival in the context of mental illness: a national cohort study*. General hospital psychiatry, 37(6), 501-506.
4. Correll, C. U., Detraux, J., De Lepeleire, J. & De Hert, M. (2015). *Effects of antipsychotics, antidepressants and mood stabilizers on risk for physical diseases in people with schizophrenia, depression and bipolar disorder*. World Psychiatry, 14(2), 119-136.
Kisely, S., Baghaie, H., Lalloo, R., Siskind, D. & Johnson, N. W. (2015). *A systematic review and meta-analysis of the association between poor oral health and severe mental illness*. Psychosomatic medicine, 77(1), 83-92.
5. Te Pou o te Whakaaro Nui (2014). *The physical health of people with a serious mental illness and/or addiction: An evidence review*. Auckland: New Zealand.

Equally Well

AOTEAROA, NEW ZEALAND - SOME KEY FACTS

Stroke

People who experience depression and anxiety are **2.3 times** more likely to have a stroke.



Co-existing issues

50% of people accessing mental health services have a substance use disorder.

70% of people accessing addiction services have a mental health need.


Tobacco smoking

People experiencing mental health issues are **2-3 times** more likely to smoke.

People experiencing substance use issues are **2-4 times** more likely to smoke.



Take action to improve the physical health of people experiencing mental health and addiction issues by:



Join the collaborative



www.tepou.co.nz/equallywell



@EquallyW

6. Lockett, H., Jury, A., Tuason, C., Lai, J. & Fergusson, D. (2018). *Comorbidities Between Mental and Physical Health Problems: An Analysis of the New Zealand Health Survey data*. New Zealand Journal of Psychology, 47(3), 5-11.

7. Todd, F.C. (2010). *Te Ariari o te Oranga: the Assessment and Management of People with Co-existing Mental Health and Substance Use Problems*. Ministry of Health, Wellington.

8. Te Pou o te Whakaaro Nui (2017). *The physical health of people with mental health conditions and/or addiction: Summary evidence update*. Auckland: New Zealand.